

STEMtaq™ DNA Polymerase Master Mix Kit



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

DNA polymerase master mix and nuclease-free water for PCR reactions

Catalog #	79009	1 Kit	100 Reactions
	79009.1	1 Kit	1000 Reactions

Product Description

STEMtaq™ DNA Polymerase Master Mix Kit, which includes master mix and nuclease-free water, reliably and consistently amplifies a wide range of PCR templates. STEMtaq™ DNA Polymerase Master Mix is a 2X concentrated, ready-to-use solution containing STEMtaq™ DNA polymerase, dNTPs, MgCl₂, and reaction buffer. STEMtaq™ DNA polymerase has 5' to 3' exonuclease activity and produces PCR fragments with a 3' A overhang.

Ordering Information

PRODUCT NAME	CATALOG #	SIZE	COMPONENTS
STEMtaq™ DNA Polymerase Master Mix Kit	79009	1 Kit - 100 Reactions	<ul style="list-style-type: none">STEMtaq™ DNA Polymerase Master Mix (79010)Nuclease-Free Water (79006)
STEMtaq™ DNA Polymerase Master Mix Kit	79009.1	1 Kit - 1000 Reactions	<ul style="list-style-type: none">STEMtaq™ DNA Polymerase Master Mix (79011)Nuclease-Free Water (79012)

Component Storage and Stability

The following components are sold as part of a kit (Catalog #79009 or 79009.1) and are not available for individual sale.

COMPONENT NAME	COMPONENT #	SIZE	STORAGE	SHELF LIFE
STEMtaq™ DNA Polymerase Master Mix*	79010	2 x 1.25 mL	Store at -20°C.	Stable until expiry date (EXP) on label.
Nuclease-Free Water	79006	2 x 1.25 mL	Store at -20°C. Alternatively, store at < 30°C.	Stable until expiry date (EXP) on label.
STEMtaq™ DNA Polymerase Master Mix*	79011	25 mL	Store at -20°C.	Stable until expiry date (EXP) on label.
Nuclease-Free Water	79012	25 mL	Store at -20°C. Alternatively, store at < 30°C.	Stable until expiry date (EXP) on label.

*Contains DNA polymerase, 400 μM dATP, 400 μM dGTP, 400 μM dCTP, 400 μM dTTP, 3 mM MgCl₂, and reaction buffer (pH 8.5).

Materials Required But Not Included

PRODUCT NAME	CATALOG #
Total DNA purification kit	e.g. Qiagen 69504
PCR tubes or plate	e.g. Corning PCR-02-C or PCR-96-C
Filtered pipette tips	e.g. 38035
Microcentrifuge tubes, 0.65 mL	e.g. 38037
Forward and reverse primers*	---
Thermocycler	---

*For assistance with primer design, visit www.stemcell.com.

Directions for Use

- Purify DNA samples using a total DNA purification kit. Store on ice.
- Thaw STEMtaq™ DNA Polymerase Master Mix and Nuclease-Free Water at room temperature (15 - 25°C).
NOTE: if not used immediately, aliquot Master Mix and store at 2 - 8°C for up to 18 weeks, or at -20°C for long-term storage. Do not exceed expiry date as indicated on label. After thawing aliquots, use immediately. Do not re-freeze.
- Vortex the Master Mix, then centrifuge briefly to collect material at the bottom of the tube.
- The following example is for preparing 50 µL of Reagent Mix. If preparing other volumes, adjust accordingly. Combine components in a microcentrifuge tube on ice as indicated in Table 1.

Table 1. Reagent Mix Components

COMPONENT	VOLUME	FINAL CONCENTRATION
STEMtaq™ DNA Polymerase Master Mix	25 µL	1X
Forward primer, 10 µM	0.5 - 5.0 µL	0.1 - 1.0 µM
Reverse primer, 10 µM	0.5 - 5.0 µL	0.1 - 1.0 µM
DNA template	1 - 5 µL	< 250 ng
Nuclease-Free Water	Variable	---
Total volume	50 µL	---

- Centrifuge the Reagent Mix for 5 seconds.
- Perform PCR in a thermocycler pre-heated to 95°C. Refer to Table 2 for recommended PCR conditions. For PCR troubleshooting, see Notes and Tips.

Table 2. Recommended PCR Cycling Conditions

STEP	TEMPERATURE	TIME
Initial denaturation	95°C	2 minutes
Annealing	~5°C below the lowest melting temperature of the primers	15 - 60 seconds
25 - 30 cycles	72 - 74°C	1 minute per kb to be amplified
	95°C	15 - 60 seconds
Final extension	72 - 74°C	5 minutes
Hold	4°C	Infinite

- Store the reaction products at 2 - 8°C for up to 24 hours or at -20°C for long-term storage.

Notes and Tips

PCR Troubleshooting

If PCR results in low amplification/no amplification of DNA, try any of the following:

- Decrease annealing temperature/increase annealing time
- Increase number of PCR cycles
- Increase concentration of primer, template, and/or polymerase
- Minimize the effect of inhibitors by diluting DNA template or using less. Alternatively, use an ethanol precipitation and wash step on DNA template prior to PCR.
- Add PCR-enhancing agents (e.g. DMSO or betaine) or a stabilizing agent such as BSA (to a final concentration of 0.16 mg/mL)

If non-specific bands are obtained:

- Increase annealing temperature
- Increase primer length to increase specificity
- Adjust annealing time:
 - If non-specific bands are longer than target, decrease annealing time
 - If non-specific bands are shorter than target, increase annealing time

Related Products

For related products, including genome editing tools, qPCR arrays, specialized cell culture and storage media, and cultureware, visit www.stemcell.com or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, STEMtaq, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. Sale of STEMtaq is covered under at least US Patent No. 6,242,235, Australian Patent No. 761757, Canadian Patent No. 2,335,153, Chinese Patent No. ZL99808861.7, Hong Kong Patent No. HK 1040262, Japanese Patent No. 3673175, European Patent No. 1088060, other patents pending, and all applicable STEMCELL Terms and Conditions of Sale. STEMtaq can be used for basic PCR experiments in research use only applications without the need to acquire a license from STEMCELL or payment of royalty fees to STEMCELL. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.