

AggreWell™400

Microwell culture plates for easy and reproducible production of embryoid bodies and spheroids



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

AggreWell™400 plates are used to generate cell aggregates such as embryoid bodies (EBs) and spheroids. Each well contains a standardized array of microwells 400 µm in size, enabling the production of large numbers of EBs and spheroids. The size of the EBs and spheroids is highly uniform and can be easily controlled by adjusting the input cell density.

If larger microwells are desired (800 µm), AggreWell™800 plates are also available (Catalog #34811/34821). AggreWell™ plates can be used to generate uniform aggregates of any cell type, which helps to ensure reproducibility of experiments.

NOTE: For all cell types, **Anti-Adherence Rinsing Solution (Catalog #07010)** is required during plate preparation steps to ensure optimal performance. Anti-Adherence Rinsing Solution prevents cell adhesion and promotes efficient EB and spheroid formation.

Storage and Stability

Store AggreWell™400 plates at room temperature (15 - 25°C) away from direct sunlight. Stable for 3 years from date of manufacture (MFG) on label.

Product Information

PRODUCT NAME	CATALOG #	SIZE	DESCRIPTION
AggreWell™400 24-well Plate	34411	1 Plate	24 wells, approximately 1200 microwells per well. Microwells are 400 µm in size.
	34415	5 Plates	
AggreWell™400 24-well Plate Starter Kit	34450	1 Kit	<ul style="list-style-type: none">2 x AggreWell™400 24-well Plates (Catalog #34411)1 x Anti-Adherence Rinsing Solution (100 mL; Catalog #07010)
AggreWell™400 6-well Plate	34421	1 Plate	6 wells, approximately 7000 microwells per well. Microwells are 400 µm in size.
	34425	5 Plates	
AggreWell™400 6-well Plate Starter Kit	34460	1 Kit	<ul style="list-style-type: none">2 x AggreWell™400 6-well Plates (Catalog #34421)1 x Anti-Adherence Rinsing Solution (100 mL; Catalog #07010)

Materials Required But Not Included

PRODUCT NAME	CATALOG #
Anti-Adherence Rinsing Solution*	07010
Trypan Blue	07050
37 µm Reversible Strainer	27215 (Small) OR 27250 (Large)
Falcon® Conical Tubes	38010 (50 mL) OR 38009 (15 mL)
Falcon® Serological Pipettes	38002 (2 mL) OR 38003 (5 mL)

*Included in AggreWell™400 24-well Plate Starter Kit (Catalog #34450) and 6-well Plate Starter Kit (Catalog #34460)

Directions for Use

NOTE: Do not expose AggreWell™ plates to organic solvents, including ethanol or isopropanol.

A. MEDIUM FOR EBs OR SPHEROIDS

Select an appropriate medium as follows:

- For generation of **EBs**:

It is essential to start with a high-quality population of undifferentiated embryonic stem (ES) or induced pluripotent stem (iPS) cells. Use an EB formation medium (e.g. AggreWell™ EB Formation Medium, Catalog #05893) supplemented with 10 μ M Y-27632 (Catalog #72302).

- For generation of **spheroids** from other cell types (including cancer spheroids):

Select an appropriate culture medium for the desired downstream application. If serum-free medium is desired, MammoCult™ Medium (Catalog #05620) may be used.

When warming medium, warm to room temperature (15 - 25°C) or 37°C as appropriate.

Please read the entire protocol before proceeding.

Use sterile techniques when performing the following protocols:

B. Preparation of AggreWell™400 Plates

C. Generation of EBs or Spheroids

D. Changing Medium in AggreWell™400 Plates

E. Harvesting EBs/Spheroids from AggreWell™400 Plates

B. PREPARATION OF AGGREWELL™400 PLATES

NOTE: For all cell types, Anti-Adherence Rinsing Solution is required during plate preparation steps to ensure optimal performance. Anti-Adherence Rinsing Solution prevents cell adhesion and promotes efficient EB and spheroid formation.

1. Warm basal medium and complete medium (see section A for further information).
2. Open AggreWell™ plates in a biosafety cabinet.

NOTE: Do not expose AggreWell™ plates to organic solvents, including ethanol or isopropanol.

3. Pre-treat wells with Anti-Adherence Rinsing Solution as described below.

- a. Add Anti-Adherence Rinsing Solution to each well to be used, as follows:

- **24-well plate:** 500 μ L
- **6-well plate:** 2 mL

- b. Centrifuge plate at 1300 x *g* for 5 minutes in a swinging bucket rotor fitted with plate holders.

NOTE: Plates must be well balanced. Prepare a balance plate using a standard plate filled with water to match the weight and position of the AggreWell™ plate.

- c. Observe plate under a microscope to ensure that bubbles have been removed from microwells. If bubbles remain trapped in any microwells, centrifuge at 1300 x *g* for an additional 5 minutes.

- d. Aspirate Anti-Adherence Rinsing Solution from the wells.

4. Rinse each well with warm basal medium, as follows:

- **24-well plate:** 2 mL
- **6-well plate:** 5 mL

Aspirate medium from the well.

5. Add warm complete medium to each well to be used, as follows:

- **24-well plate:** 1 mL
- **6-well plate:** 2.5 mL

C. GENERATION OF EBs OR SPHEROIDS

1. Prepare a single-cell suspension in desired medium (see section A).
2. Perform a cell count using Trypan Blue and a hemocytometer to determine the viable cell concentration.

3. Refer to Table 1 to determine the number of cells required per well to achieve the desired number of cells per microwell. Alternatively, calculate using the formula for either a 24-well or a 6-well plate, as follows:

24-well plate: Required number of cells per well = Desired number of cells per microwell x 1200 microwells per well

6-well plate: Required number of cells per well = Desired number of cells per microwell x 7000 microwells per well

Table 1. Required Number of Cells per Well for AggreWell™400 Plates

24-WELL PLATES		6-WELL PLATES	
Desired number of cells per microwell*	Required number of cells per well	Desired number of cells per microwell*	Required number of cells per well
50	6×10^4	50	3.5×10^5
100	1.2×10^5	100	7.0×10^5
200	2.4×10^5	200	1.4×10^6
500	6×10^5	500	3.5×10^6
1000	1.2×10^6	1000	7.0×10^6
2000	2.4×10^6	2000	1.4×10^7

*The recommended range is 50 - 3000 cells per microwell.

NOTE: For most cell types, the number of cells per microwell will equal the number of cells per spheroid (i.e. 100% incorporation); for some cell lines or cell types, incorporation may be less than 100%. For EBs, not all ES or iPS cells will be incorporated into the aggregate.

4. Adjust the concentration of the single-cell suspension and add a sufficient volume to each well to achieve the desired cell number as per Table 1.

NOTE: Avoid performing multiple dispensing steps from a single aspiration of the cell suspension as this may reduce the accuracy of seeding numbers in each well.

5. Add complete medium to each well to achieve a final volume as follows:

- **24-well plate:** 2 mL/well
- **6-well plate:** 5 mL/well

6. Prepare a centrifuge balance plate using a standard plate filled with water to match the weight and position of the AggreWell™ plate.
7. Pipette cells up and down gently several times to ensure even distribution of cells throughout the well. Be careful not to introduce bubbles into the microwells.
8. Immediately centrifuge the AggreWell™ plate at $100 \times g$ for 3 minutes to capture cells in the microwells, using the balance plate prepared in step 6.
9. Observe plate under a microscope to verify that cells are evenly distributed among the microwells.
10. Incubate the plate at 37°C with 5% CO_2 and 95% humidity for 24 hours. Observe the cells under a microscope.

NOTE: Many cell lines form EBs/spheroids within 24 hours, but some may require a longer incubation time (up to 48 hours) for optimal EB/spheroid formation.

D. CHANGING MEDIUM IN AGGREWELL™400 PLATES

Some applications of EBs or spheroids may require continuous culture in AggreWell™ plates, including medium changes. For best results, use a serological pipette (2 mL or 5 mL) for medium changes.

Perform medium changes as described below:

1. Warm complete medium.
2. Perform a 50 - 75% medium change as follows:

24-well plate

- a. Slowly remove 1 - 1.5 mL of medium from each well.
- b. Replace with 1 - 1.5 mL of fresh complete medium by slowly pipetting down the wall of the well. Slowly dispensing the medium helps to prevent displacement of EBs/spheroids from the microwells.

6-well plate

- a. Slowly remove 2.5 - 3.75 mL of medium from each well.
- b. Replace with 2.5 - 3.75 mL of fresh complete medium by slowly pipetting down the wall of the well. Slowly dispensing the medium helps to prevent displacement of EBs/spheroids from the microwells.

For culturing cells in AggreWell™400 plates, refer to the appropriate culture protocol for further instructions. For harvesting EBs/spheroids from AggreWell™400 plates, proceed to section E.

E. HARVESTING EBs/SPHEROIDS FROM AGGREWELL™400 PLATES

1. Warm basal medium and complete medium.
2. Using a serological pipette:
 - a. Remove approximately half of the culture medium from the well.
 - b. Dispense the medium firmly back onto the surface of the plate to dislodge the EBs/spheroids from the microwells. Do not triturate.
3. Select the appropriate strainer and conical tube for separation of EBs/spheroids from single cells:

24-well plate

- For harvesting from a single well, use 37 µm Reversible Strainer, Small and a 15 mL conical tube
- For harvesting from multiple wells, use 37 µm Reversible Strainer, Large and a 50 mL conical tube

6-well plate

- For harvesting from a single well, use 37 µm Reversible Strainer, Large and a 50 mL conical tube

4. Place strainer on top of the tube with the arrow pointed upward.
5. Gently aspirate the dislodged EBs/spheroids (from step 2). Pass the EB/spheroid suspension through the strainer.
NOTE: The aggregates will remain on the filter; any unincorporated single cells will flow through.
6. Using a serological pipette, dispense 1 mL (**24-well plate**) or 3 mL (**6-well plate**) of warm basal medium across the entire surface of the well to dislodge any remaining EBs/spheroids. Collect wash and pass over the strainer used in step 5. Repeat this wash step 3 times.
7. Invert the strainer, and place over a new conical tube of the same size. Collect the EBs/spheroids by washing with 2 - 5 mL of complete medium per well harvested.
8. Observe the AggreWell™ plate under a microscope to ensure that all EBs/spheroids have been removed from the wells. Repeat wash if necessary (steps 6 - 7).
9. OPTIONAL: Count the EBs/spheroids to determine actual yield, as described below.

Using a serological pipette:

- a. Pipette the EB/spheroid suspension up and down 2 - 3 times to ensure even distribution.
- b. Pipette 50 µL of the EB/spheroid suspension into a flat-bottom 96-well plate. Count at 20 - 100X magnification.
- c. Calculate EB or spheroid yield as follows:

$$\text{Total number of EBs or spheroids} = \frac{\text{EB or spheroid count (in 50 } \mu\text{L)}}{50 \mu\text{L}} \times \text{Volume of EB or spheroid suspension (}\mu\text{L)}$$

NOTE: The expected yield is approximately 1200 EBs/spheroids for 1 well of a 24-well plate, or 7000 EBs/spheroids for 1 well of a 6-well plate.

10. Harvested EBs/spheroids are now ready for downstream applications such as suspension culture, directed differentiation (EBs), drug screening and toxicity assays, and analysis.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, AggreWell, and MammoCult are trademarks of STEMCELL Technologies Canada Inc. Falcon is a registered trademark of Corning Incorporated. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.