

Cytokines

Human Recombinant IP-10 (CXCL10)

Interferon gamma-inducible protein 10

Catalog # 78165
78165.1

10 µg
50 µg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Interferon gamma-inducible protein (IP) 10 or CXCL10 is a member of the CXC chemokine family. It binds CXCR3 activating ERK1/2, p38/MAPK, JNK, and PI3-kinase/AKT signaling pathways, inducing intracellular calcium influx, DNA synthesis, cell proliferation, and chemotaxis. IP-10 regulates innate and adaptive immune responses by affecting the function of activated T cells, natural killer cells, inflammatory dendritic cells, macrophages, and B cells. IP-10 is produced by leukocytes, activated neutrophils, eosinophils, monocytes, epithelial cells, endothelial cells, fibroblasts, and keratinocytes in response to IFN-gamma. IP-10 has been implicated in a wide range of diseases, including infectious diseases and chronic inflammatory and autoimmune diseases, as well as in tumor formation (Liu et al.).

Product Information

Alternative Names: 10 kDa interferon-gamma-induced protein, C7, crg-2, Chemokine (C-X-C motif) ligand 10, CXCL10, Gamma-IP-10, gIP-10, IFI10, INP10, mob1, SCYB10, Small inducible cytokine B10

Accession Number: P02778

Amino Acid Sequence: MVPLSRTVRC TCISISNQPV NPRSLEKLEI IPASQFCPRV EIIATMKKKG EKRCNLNPEK AIKNLLKAVS KEMSKRSP

Predicted Molecular Mass: 8.8 kDa

Species: Human

Cross Reactivity: Reported to be species-specific

Formulation: Lyophilized after dialysis against Tris buffer, pH 8.0.

Source: E. coli

Specifications

Activity: The specific activity is $\geq 5.0 \times 10^3$ units/mg ($EC_{50} \leq 0.2 \mu\text{g/mL}$) as determined by a Ca^{2+} mobilization assay using CHO-K1 cells expressing human $G\alpha 15$ and mouse CXCR3.

Purity: $\geq 95\%$

Endotoxin Level: Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 0.2 EU/ μg protein.

Preparation and Storage


Storage: Store at -80°C .

Stability: Stable as supplied for 12 months from date of receipt.

Preparation: Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at $2 - 8^\circ\text{C}$ for more than 2 weeks or at -20°C to -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant IP-10 (CXCL10) was tested using a Ca²⁺ mobilization assay in CHO-K1 stably expressing Gα15 and CXCR3. Calcium mobilization was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which calcium mobilization is at 50% of maximum. The EC₅₀ in the example above is less than 0.2 µg/mL.

(B) 2 µg of Human Recombinant IP-10 (CXCL10) was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant IP-10 (CXCL10) has a predicted molecular mass of 8.8 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Liu M et al. (2011) The emerging role of CXCL10 in cancer (Review). *Oncol Lett* 2(4): 583–9.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.