

Cytokines

Human Recombinant FGF-5



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Fibroblast growth factor 5

Catalog # 78182
78182.1

50 µg
100 µg

Product Description

Fibroblast growth factor 5 (FGF-5) is a secreted, heparin-binding member of the FGF subfamily. FGFs possess broad mitogenic and cell survival activities and are expressed during embryonic development. FGF-5 is expressed in the mesenchyme, skeletal muscles, central nervous systems, and hair follicles. FGF-5 promotes cell differentiation and proliferation by binding FGF receptor 1 and FGF receptor 2 (FGFR1 and FGFR2, respectively). FGF-5 plays an important regulatory role in skeletal muscle development. FGF-5 has also been identified in neurons of the limbic system, especially those of the olfactory bulb and pyramidal cells of the hippocampus (Haub & Goldfarb).

Product Information

Alternative Names: Fibroblast growth factor 5, HBGF-5, Heparin-binding growth factor 5, Smag-82
Accession Number: P12034
Amino Acid Sequence: MAWAHGEKRL APKGQPGPAA TDRNPIGSSS RQSSSSAMSS SSASSSPAAS LGSQGSGLLEQ SSFQWSPSGR RTGSLYCRVG IGFHLQIYPD GKVNGSHEAN MLSVLEIFAV SQGIVGIRGV FSNKFLAMSK KGKLHASAKF TDDCKFRERF QENSYNTYAS AIHRTEKTGR EWYVALNKRK KAKRGCSPRV KPQHISTHFL PRFKQSEQPE LSFVTVPEK KNPPSPIKSK IPLSAPRKNT NSVKYRLKFR FG
Predicted Molecular Mass: 27.7 kDa
Species: Human
Cross Reactivity: Reported to be species-specific
Formulation: Lyophilized from a sterile-filtered solution containing sodium phosphate and sodium chloride, pH 7.5.
Source: E. coli

Specifications


Activity: The specific activity is $\geq 1.0 \times 10^5$ units/mg ($EC_{50} \leq 10$ ng/mL) as determined by a cell proliferation assay using NR6R-3T3 cells in the presence of 1 µg heparin.
Purity: $\geq 95\%$
Endotoxin Level: Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage: Store at -20°C to -80°C.
Stability: Stable as supplied for 12 months from date of receipt.
Preparation: Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant FGF-5 was tested by its ability to promote proliferation of NR6R-3T3 cells in the presence of 1 μ g heparin. Cell proliferation was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC₅₀ in the example above is 3.11 ng/mL.

(B) 1 μ g of Human Recombinant FGF-5 was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant FGF-5 has a predicted molecular mass of 27.7 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Haub O & Goldfarb M. (1991) Expression of the fibroblast growth factor-5 gene in the mouse embryo. *Development* 112(2): 397–406.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.