

Small Molecules

LDN193189

BMP pathway inhibitor; Inhibits ALK1, ALK2, ALK3, and ALK6

Catalog # 72147
72149

10 mg
50 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

LDN193189 is a potent inhibitor of the bone morphogenetic (BMP) pathway, inhibiting ALK1, ALK2, ALK3, and ALK6 (IC_{50} = 0.8, 0.8, 5.3, and 16.7 nM respectively; Sanvitale et al.). It is a derivative of Dorsomorphin (Catalog #72102) that is typically used at approximately 100-fold lower concentrations (Sanvitale et al.; Vogt et al.). This product is supplied as the dihydrochloride salt of the molecule and it is the most soluble version of LDN193189.

Molecular Name:	LDN193189 (Dihydrochloride)
Alternative Names:	DM-3189
CAS Number:	1435934-00-1
Chemical Formula:	$C_{25}H_{22}N_6 \cdot 2HCl$
Molecular Weight:	479.4 g/mol
Purity:	> 98%
Chemical Name:	4-[6-[4-(1-piperazinyl)phenyl]pyrazolo[1,5-a]pyrimidin-3-yl]-quinoline dihydrochloride
Structure:	



Properties

Physical Appearance:	Orange solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For long-term storage store with a desiccant. Stable as supplied for 12 months from date of receipt.
Solubility:	· Water \leq 50 mM · DMSO \leq 20 mM For example, to prepare a 10 mM stock solution in water, resuspend 10 mg in 2.09 mL of water. NOTE: This is based on a molecular weight (MW) of 479.4 g/mol. MW may vary due to water content of the molecule. For batch-specific MW, request a Certificate of Analysis at techsupport@stemcell.com.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

DIFFERENTIATION

- Promotes differentiation of neural progenitor cells from human pluripotent stem cells (Chambers et al.; Kriks et al.).
- Promotes differentiation of neural crest cells from human pluripotent stem cells (Kreitzer et al.).
- Promotes differentiation of anterior foregut endoderm from human and mouse pluripotent stem cell-derived definitive endoderm (Kearns et al.).
- Promotes differentiation of sensory epithelial cells of the inner ear from mouse embryonic stem cells (Koehler et al.).

CANCER RESEARCH

- Inhibits prostate and breast cancer tumor growth (Balboni et al.; Lee et al.).
- Prevents ovarian cancer cell proliferation (Tsai et al.).

References

- Balboni AL et al. (2013) Δ Np63 α -Mediated activation of bone morphogenetic protein signaling governs stem cell activity and plasticity in normal and malignant mammary epithelial cells. *Cancer Res* 73(2).
- Chambers SM et al. (2012) Combined small-molecule inhibition accelerates developmental timing and converts human pluripotent stem cells into nociceptors. *Nat Biotechnol* 30(7): 715–20.
- Kearns NA et al. (2013) Generation of organized anterior foregut epithelia from pluripotent stem cells using small molecules. *Stem Cell Res* 11(3): 1003–12.
- Koehler KR et al. (2013) Generation of inner ear sensory epithelia from pluripotent stem cells in 3D culture. *Nature* 500(7461): 217–21.
- Kreitzer FR et al. (2013) A robust method to derive functional neural crest cells from human pluripotent stem cells. *Am J Stem Cells* 2(2): 119–31.
- Kriks S et al. (2011) Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease. *Nature* 480(7378): 547–51.
- Lee C-T et al. (2016) CYP3A5 mediates effects of cocaine on human neocortico genesis: studies using an in vitro 3D self-organized hPSC model with a single cortex-like unit. *Neuropsychopharmacology*.
- Lee Y-C et al. (2011) BMP4 promotes prostate tumor growth in bone through osteogenesis. *Cancer Res* 71(15): 5194–203.
- Sanvitale CE et al. (2013) A new class of small molecule inhibitor of BMP signaling. *PLoS One* 8(4): e62721.
- Tsai C-L et al. (2012) Secreted stress-induced phosphoprotein 1 activates the ALK2-SMAD signaling pathways and promotes cell proliferation of ovarian cancer cells. *Cell Rep* 2(2): 283–93.
- Vogt J et al. (2011) The specificities of small molecule inhibitors of the TGF β and BMP pathways. *Cell Signal* 23(11): 1831–42.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. Sold under license of at least US Patent Nos. 8,507,501 and 9,045,484, and any foreign counterparts thereof. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.