

Cytokines

Human Recombinant RANKL



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Receptor activator of nuclear factor
kappa-B ligand

Catalog #	78214	10 µg
	78214.1	100 µg
	78214.2	500 µg
	78214.3	1000 µg

Product Description

Receptor activator of nuclear factor kappa-B ligand (RANKL) is a member of the tumor necrosis factor (TNF) superfamily (Anderson et al.). Cytokines in the TNF superfamily are involved in a variety of long-term cellular activities, such as differentiation, proliferation, and cell death (MacEwan). RANKL is a type II homotrimeric transmembrane protein expressed in both a membrane-bound and secreted form (Ikeda et al.). RANKL binds to the receptor activator of nuclear factor kappa-B (RANK). Upon binding to its receptor, RANKL activates the AKT signaling pathway (Moon et al.). Osteoprotegerin (OPG) may also bind RANKL, and this binding competes with RANKL-RANK binding (Lacey et al.). RANKL is involved in osteoclastogenesis (Lacey et al.; Yasuda et al.) and T cell activation (Wong et al.).

Product Information

Alternative Names:	CD254, hRANKL2, ODF, OPGL, OPTB2, Osteoclast differentiation factor, soluble Receptor activator of NF-κB ligand, sOdf, TNF-related activation-induced cytokine, TNFSF11, TNF superfamily member 11, TNLG6B, Tumor necrosis factor superfamily member 11, TRANCE
Accession Number:	O14788
Amino Acid Sequence:	EKAMVDGSWL DLAKRSKLEA QPFAHLTINA TDIPSGSHKV SLSSWYHDRG WAKISNMTFS NGKLIVNQDG FYYLYANICF RHHETSGDLA TEYLQLMVYV TKTSIKIPSS HTLMKGGSTK YWSGNSEFHF YSINVGGFFK LRSGEIEISIE VSNPSLLDPD QDATYFGAFK VRDID
Predicted Molecular Mass:	19.7 kDa
Species:	Human
Cross Reactivity:	Mouse
Formulation:	Lyophilized from a sterile-filtered solution containing sodium phosphate, pH 7.5.
Source:	E. coli

Specifications


Activity:	The specific activity is $\geq 2.0 \times 10^4$ units/mg ($EC_{50} \leq 50$ ng/mL) as determined by embryonic alkaline phosphatase production induced in RAW-Blue™ macrophage reporter cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage:	Store at -20°C to -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -20°C to -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant RANKL was tested by its ability to induce production of embryonic alkaline phosphatase in RAW-Blue™ macrophage reporter cells. Alkaline phosphatase production was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which alkaline phosphatase activity is at 50% of maximum. The EC₅₀ in the above example is 38.8 ng/mL.

(B) Human Recombinant RANKL was resolved with SDS-PAGE under reducing (+) conditions and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant RANKL has a predicted molecular mass of 19.7 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Anderson DM et al. (1997) A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature* 390(6656): 175–9.
- Ikeda T et al. (2001) Determination of three isoforms of the receptor activator of nuclear factor-kappaB ligand and their differential expression in bone and thymus. *Endocrinology* 142(4): 1419–26.
- Lacey DL et al. (1998) Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 93(2): 165–76.
- MacEwan DJ. (2002) TNF ligands and receptors—a matter of life and death. *Br J Pharmacol* 135(4): 855–75.
- Moon JB et al. (2012) Akt induces osteoclast differentiation through regulating the GSK3β/NFATc1 signaling cascade. *J Immunol* 188(1): 163–9.
- Wong BR et al. (1997) TRANCE is a novel ligand of the tumor necrosis factor receptor family that activates c-Jun N-terminal kinase in T cells. *J Biol Chem* 272(40): 25190–4.
- Yasuda H et al. (1998) Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci USA* 95(7): 3597–602.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2019 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. RAW-Blue is a trademark of InvivoGen. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.