


SnapShot: Directed Differentiation of ESCs and iPSCs

Cell

Luis A. Williams, Brandi N. Davis-Dusenberry, and Kevin C. Eggan
HHMI, Harvard University, Cambridge, MA 02138, USA

This SnapShot was previously published in Cell 149, May 25, 2012 ©2012 Elsevier Inc. DOI 10.1016/j.cell.2012.05.015



STEMCELL Technologies is committed to making sure your research works. As Scientists Helping Scientists, we support our customers by creating novel products with consistent, unfailing quality and by providing unparalleled technical support.

mTeSR™ Plus feeder-free medium for hPSC maintenance offers enhanced culture stability for consistent and high quality cultures upstream of your differentiation workflow. To find out more and to explore a range of feeder-free culture media, visit www.tesr.com.

Enjoy a streamlined hPSC-derived differentiation workflow with mTeSR™ Plus and STEMdiff™, a complete system for reproducible differentiation to specialized cell types and organoids. Learn more at www.stemcell.com/STEMdiff.

For the cytokines and small molecules shown above, explore online:

- www.stemcell.com/PSCcytokines
- www.stemcell.com/PSCsmallmolecules

STEMCELL™
TECHNOLOGIES
Scientists Helping Scientists™ | WWW.STEMCELL.COM